

# Sun and shade leaf variability in *Liquidambar chinensis* and *Liquidambar formosana* (Altingiaceae): Implications for palaeobotany

Maslova N., Karasev E., Kodrul T., Spicer R., Volkova L., Spicer T., Jin J., Liu X.  
*Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia*

---

## Abstract

© 2018 The Linnean Society of London. Many factors influence leaf anatomy and morphology in the crown of a tree, particularly those resulting from microclimatic differences between the periphery and the interior of the crown. These influences can be so strong that single species can produce different leaf forms in which shade and sun leaves exhibit consistently distinctive morphological and epidermal character sets. Here we show, using *Liquidambar* as a model system, that the principal morphological characters for distinguishing shade and sun leaves in two modern *Liquidambar* spp. with different lamina types (entire in *L. chinensis* and lobate in *L. formosana*) are the leaf lamina length to width ratio, the degree of development of venation networks, tooth size and tooth shape. The main epidermal characters are ordinary cell size and anticlinal wall outlines. Many fossils, however, are only preserved as impressions and morphological characters alone have been used to distinguish shade and sun leaf morphotypes. To evaluate the utility of our approach, populations of fossil *Liquidambar* leaves from the Eocene of southern China, preserved only as impressions, were categorized into sun and shade morphotypes. Recognition that sun and shade leaf morphological diversity exists in fossil populations will enable palaeobotanists to identify more reliably foliar polymorphisms that would otherwise be used to describe, incorrectly, different species.

<http://dx.doi.org/10.1093/botlinnean/boy047>

---

## Keywords

Epidermal characters, Fossil leaves, Leaf morphology, Statistics

## References

- [1] Balsamo RA, Bauer AM, Davis SD, Rice BM. 2003. Leaf biomechanics, morphology, and anatomy of the deciduous mesophyte *Prunus serrulata* (Rosaceae) and the evergreen sclerophyllous shrub *Heteromeles arbutifolia* (Rosaceae). *American Journal of Botany* 90: 72-77
- [2] Barbacka M, van Konijnenburg-van Cittert JHA. 1998. Sun and shade leaves in two Jurassic species of Pteridosperms. *Review of Palaeobotany and Palynology* 103: 209-221
- [3] Bentham G, Hooker JD. 1865. *Genera plantarum*. London: Reeve and Williams & Norgate
- [4] Blume CL. 1828. *Flora Javae nec non insularum adjacentium*. Brussels: J. Frank
- [5] Brown RW. 1933. A Cretaceous sweet gum. *Botanical Gazette* 94: 611-615
- [6] Chaney RW, Axelrod DI. 1959. Miocene floras of the Columbia Plateau. Part II, systematic considerations. *Carnegie Institution of Washington Publication* 617: 135-229

- [7] Cooke R. 2005. The history of mathematics: a brief course. 2nd edn. Hoboken: Wiley-Interscience
- [8] Ellis B, Daly DC, Hickey LJ, Johnson KR, Mitchell JD, Wilf P, Wing SL. 2009. Manual of leaf architecture. Ithaca: Cornell University Press
- [9] Endo S, Morita H. 1932. Notes on the genera *Comptoniophyllum* and *Liquidambar*. Science reports of the Tohoku Imperial University. 2nd series, Geology 15: 41-53
- [10] Givnish TJ. 1988. Adaptation to sun and shade: a whole-plant perspective. Australian Journal of Plant Physiology 15: 63-92
- [11] Golovneva LB. 2004. The Late Cretaceous flora of Siberia. Doctoral Dissertation in Biology, Komarov Botanical Institute
- [12] Greenwood JA, Sandomire MM. 1950. Sample size required for estimating the standard deviation as a per cent of its true value. Journal of the American Statistical Association 45: 257-260
- [13] Guignard G, Bóka K, Barbacka M. 2001. Sun and shade leaves? Cuticle ultrastructure of Jurassic *Komlopteris nordenskiöldii* (Nathorst) Barbacka. Review of Palaeobotany and Palynology 114: 191-208
- [14] Herrick JD, Maherali H, Thomas RB. 2004. Reduced stomatal conductance in sweetgum (*Liquidambar styraciflua*) sustained over long-term CO enrichment. New Phytologist 162: 387-396
- [15] Hu HH, Chaney RW. 1940. A Miocene flora from Shantung province, China: Part I. Introduction and systematic considerations. Washington: Carnegie Institution of Washington
- [16] Huzioka K. 1972. The Tertiary floras of Korea. Journal of the Mining College, Akita University. Ser. A, Mining Geology 5: 1-83
- [17] Huzioka K, Uemura K. 1979. The *Comptonia*-*Liquidambar* forest during middle Miocene Daijima age in Japan. Report of the Research Institute of Underground Resources, Mining College, Akita University 45: 37-50
- [18] Ickert-Bond SM, Pigg KB, Wen J. 2005. Comparative infructescence morphology in *Liquidambar* (Altingiaceae) and its evolutionary significance. American Journal of Botany 92: 1234-1255
- [19] Ickert-Bond SM, Pigg KB, Wen J. 2007. Comparative infructescence morphology in *Altingia* (Altingiaceae) and discordance between morphological and molecular phylogenies. American Journal of Botany 94: 1094-1115
- [20] Ickert-Bond SM, Wen J. 2006. Phylogeny and biogeography of Altingiaceae: evidence from combined analysis of five non-coding chloroplast regions. Molecular Phylogenetics and Evolution 39: 512-528
- [21] Ickert-Bond S, Wen J. 2013. A taxonomic synopsis of Altingiaceae with nine new combinations. PhytoKeys 31: 21-61
- [22] Knowlton FH. 1902. Fossil flora of the John Day basin, Oregon. Washington: Government Printing Office
- [23] Kouwenberg LLR, Kürschner WM, McElwain JC. 2007. Stomatal frequency change over altitudinal gradients: prospects for paleoaltimetry. Reviews in Mineralogy and Geochemistry 66: 215-241
- [24] Krzywinski M, Altman N. 2014. Visualizing samples with box plots. Nature Methods 11: 119-120
- [25] Kürschner WM. 1997. The anatomical diversity of recent and fossil leaves of the durmast oak (*Quercus petraea* Lieblein/*Q. pseudocastanea* Goepfert)-implications for their use as biosensors of palaeoatmospheric CO levels. Review of Palaeobotany and Palynology 96: 1-30
- [26] Larcher W. 1976. ökologie der Pflanzen, 2nd edn. Stuttgart: Ulmer
- [27] Leaf Architecture Working Group. 1999. Manual of leaf architecture: morphological description and categorization of dicotyledonous and net-veined monocotyledonous angiosperms. Washington: Leaf Architecture Working Group
- [28] Lindley J. 1836. A natural system of botany. London: Longman. MacGinitie HD. 1941. A Middle Eocene flora from the central Sierra Nevada. Washington: Carnegie Institution
- [29] Maslova NP. 1995. *Liquidambar* L. from the Cenozoic of eastern Asia. Paleontological Journal 29 (1A): 145-158
- [30] Maslova NP, Gordenko NV, Volkova LD. 2008. Structural epidermal features of leaves of *Platanus acerifolia* Willd. (Platanaceae) and significance of cuticular analysis for the determination of Cretaceous *Platanus*-like leaves. Botanicheskii Zhurnal 93: 982-995 [In Russian]
- [31] Maslova NP, Kodrul TM, Song Y, Volkova LD, Jin J. 2015. *Liquidambar maomingensis* sp. nov. (Altingiaceae) from the late Eocene of South China. American Journal of Botany 102: 1356-1370
- [32] Maslova NP, Shilin PV. 2011. The new species *Ettingshausenia sarbaensis* (Angiospermae) from the Cenomanian-Turonian of western Kazakhstan in light of the problem of classification of dispersed Cretaceous *Platanus*-like leaves. Paleontological Journal 45: 459-473
- [33] Maslova NP, Volkova LD, Gordenko NV. 2008. Morphological variability of leaves of *Platanus acerifolia* Willd. (Platanaceae) and approaches to the determination of dispersed Cretaceous *Platanus*-like leaves. Botanicheskii Zhurnal 93: 825
- [34] Meyer HW, Manchester SR. 1997. The Oligocene Bridge Creek flora of the John Day Formation, Oregon. Berkeley, London: University of California Press
- [35] Oken L. 1841. Allgemeine Naturgeschichte für alle Stände. Stuttgart: Hoffmann'sche Verlags-Buchhandlung

- [36] Onoe T. 1974. A Middle Miocene flora from Oguni-Machi, Yamagata prefecture, Japan. Reports, Geological Survey of Japan 253: 1-64
- [37] Ozaki K. 1991. Late Miocene and Pliocene floras in central Honshu, Japan. Yokohama: Kanagawa Prefectural Museum
- [38] Pandey S, Nagar PK. 2002. Leaf surface wetness and morphological characteristics of *Valeriana jatamansi* grown under open and shade habitats. *Biologia Plantarum* 45: 291-294
- [39] Pigg KB, Ickert-Bond SM, Wen J. 2004. Anatomically preserved *Liquidambar* (Altingiaceae) from the middle Miocene of Yakima Canyon, Washington State, USA, and its biogeographic implications. *American Journal of Botany* 91: 499-509
- [40] Poole I, Kürschner WM. 1999. Stomatal density and index: the practice. In: Jones TP, Rowe NP, eds. *Fossil plants and spores: modern techniques*. London: Geological Society, 257-260
- [41] Poole I, Weyers JDB, Lawson T, Raven JA. 1996. Variations in stomatal density and index: implications for palaeoclimatic reconstructions. *Plant, Cell and Environment* 19: 705-712
- [42] Pigliucci M. 2003. Phenotypic integration: studying the ecology and evolution of complex phenotypes. *Ecology Letters* 6: 265-272
- [43] R Core Team. 2014. R: a language and environment for statistical computing. Available at: <https://www.r-project.org>
- [44] Rozendaal DMA, Hurtado VH, Poorter L. 2006. Plasticity in leaf traits of 38 tropical tree species in response to light; relationships with light demand and adult stature. *Functional Ecology* 20: 207-216
- [45] RStudio Team. 2015. RStudio: integrated development for R. Boston: RStudio, Inc. Available at: <http://www.rstudio.com>
- [46] Rubio de Casas R, Vargas P, Pérez-Corona E, Manrique E, García-Verdugo C, Balaguer L. 2011. Sun and shade leaves of *Olea europaea* respond differently to plant size, light availability and genetic variation. *Functional Ecology* 25: 802-812
- [47] Sack L, Melcher PJ, Liu WH, Middleton E, Pardee T. 2006. How strong is intracanopy leaf plasticity in temperate deciduous trees? *American Journal of Botany* 93: 829-839
- [48] Samsonov SK. 1964. New data on Upper Cretaceous flora of Karatau. In: *Biostratigraphy and paleogeography of Meso-Cenozoic oil and gas-bearing regions of the south-east of the USSR*. Moscow: Nauka, 31-35. [In Russian]
- [49] Shi S, Huang Y, Zhong Y, Du Y, Zhang Q, Chang H, Boufford DE. 2001. Phylogeny of the Altingiaceae based on cpDNA matK, PY-IGS and nrDNA ITS sequences. *Plant Systematics and Evolution* 230: 13-24
- [50] Smith RF. 1967. The leaf dimorphism of *Liquidambar styraciflua* L. *The American Midland Naturalist* 77: 42-50
- [51] Stults DZ, Axsmith BJ. 2011. Filling the gaps in the Neogene plant fossil record of eastern North America: new data from the Pliocene of Alabama. *Review of Palaeobotany and Palynology* 167: 1-9
- [52] Sun B, Dilcher DL, Beerling DJ, Zhang C, Yan D, Kowalski E. 2003. Variation in *Ginkgo biloba* L. leaf characters across a climatic gradient in China. *Proceedings of the National Academy of Sciences of the United States of America* 100: 7141-7146
- [53] Suzuki K. 1961. The important and characteristic Pliocene and Miocene species of plants from the southern part of the Tohoku District, Japan. Fukushima University, Faculty of Art and Science
- [54] Tsukaya H. 2005. Leaf shape: genetic controls and environmental factors. *The International Journal of Developmental Biology* 49: 547-555
- [55] Uemura K. 1983. Late Neogene *Liquidambar* (Hamamelidaceae) from the southern part of northeast Honshu, Japan. *Memoirs of the National Science Museum of Tokyo* 16: 25-36
- [56] Witham E, Marchiano M, Reynolds D. 2014. Relationship between canopy location and tensile strength of leave of *Heteromeles arbutifolia* in Malibu, CA. All Undergraduate Student Research: Paper 146. Available at: <http://digitalcommons.pepperdine.edu/sturesearch/146>
- [57] Wolfe JA. 1993. A method of obtaining climatic parameters from leaf assemblages. *U. S. Geological Survey Bulletin* 2040: 1-71
- [58] Wolfe JA, Tanai T. 1980. The Miocene Seldovia Point flora from the Kenai Group, Alaska. *Professional Paper* 1105
- [59] Wu J, Sun B, Liu Y-S, Xie S, Lin Z. 2009. A new species of *Exbucklandia* (Hamamelidaceae) from the Pliocene of China and its paleoclimatic significance. *Review of Palaeobotany and Palynology* 155: 32-41
- [60] Xiao L, Sun B, Li X, Ren W, Jia H. 2011. Anatomical variations of living and fossil *Liquidambar* leaves: a proxy for paleoenvironmental reconstruction. *Science China Earth Sciences* 54: 493-508
- [61] Xiao L, Yang H, Sun B, Li X, Guo J. 2013. Stable isotope compositions of recent and fossil sun/shade leaves and implications for palaeoenvironmental reconstruction. *Review of Palaeobotany and Palynology* 190: 75-84
- [62] Xu F, Guo W, Xu W, Wei Y, Wang R. 2009. Leaf morphology correlates with water and light availability: what consequences for simple and compound leaves? *Progress in Natural Science* 19: 1789-1798

- [63] Yang J, Spicer RA, Spicer TEV, Arens NC, Jacques FMB, Su T, Kennedy EM, Herman AB, Steart DC, Srivastava G, Mehrotra RC, Valdes PJ, Mehrotra NC, Zhou ZK, Lai J-S. 2015. Leaf form-climate relationships on the global stage: an ensemble of characters. *Global Ecology and Biogeography* 24: 1113-1125
- [64] Zалensky VR. 1904. Materials to numerical anatomy of various leaves in the same plants. *Izvestiya Kievskogo Politekhnikheskogo Instituta* 4: 1-209. [In Russian]
- [65] Zhang Z, Lu A. 1995. Hamamelidaceae: geographic distribution, fossil history and origin. *Chinese Science Abstracts Series B* 33: 313-339